	P
	C
	S
	T
	C
	P
	E
	R
	Α
	T
	I
	ν
	Е
	I
	N
	S
	T
	R
	ı
	C
	T
	ı
	C
	N
	S

DIET First Hour (1) — Patient may slowly begin with CLEAR liquids (water, Gatorade, apple juice, etc). Second Hour (2) — Patient may proceed to soft foods. Third Hour (3) — If patient is doing well with liquids and softs, may slowly progress to regular diet. Avoid meat, dairy, and greasy foods. In addition, the dentist may prescribe certain dietary restrictions. PAIN & DISCOMFORT It is not uncommon for the patient to experience pain after his/her dental procedure. The anesthesia (numbing) to help the patient deal with potential pain. These medications typically wear off after a couple of hours. The following medications may be recommended for post-operative pain control. If the patient is experiencing persistent pain after taking recommended for post-operative pain control. If the patient is experiencing persistent pain after taking recommended medications, please contact the DENTIST. Tylenol/Acetaminophen (dose per package instructions) Immediately				
First Hour (1) — Patient may slowly begin with CLEAR liquids (water, Gatorade, apple juice, etc). Second Hour (2) — Patient may proceed to soft foods. Third Hour (3) — If patient is doing well with liquids and softs, may slowly progress to regular diet. Avoid meat, dairy, and greasy foods. In addition, the dentist may prescribe certain dietary restrictions. PAIN & DISCOMFORT It is not uncommon for the patient to experience pain after his/her dental procedure. The anesthesiologist may have administered IV pain medications, and the dentist may have administered local anesthesia (numbing) to help the patient deal with potential pain. These medications typically wear off after a couple of hours. The following medications may be recommended medications, splease contact the DENTIST. Tylenol/Acetaminophen (dose per package instructions) Immediately	PATIENT NAME:	Date:		
It is not uncommon for the patient to experience pain after his/her dental procedure. The anesthesiologist may have administered IV pain medications, and the dentist may have administered local anesthesia (numbing) to help the patient deal with potential pain. These medications typically wear off after a couple of hours. The following medications may be recommended for post-operative pain control. If the patient is experiencing persistent pain after taking recommended medications, please contact the DENTIST. Tylenol/Acctaminophen (dose per package instructions)	First Hour (1) – Patient may slowly begin with CLEAR liquids (water, Gatorade, apple juice, etc). Second Hour (2) – Patient may proceed to soft foods. Third Hour (3) – If patient is doing well with liquids and softs, may slowly progress to regular diet.			
OR - Ibuprofen/Advil/Motrin (dose per package instructions) Immediately Begin atam/pm NAUSEA & VOMITING Occasionally nausea and vomiting may occur following anesthesia. Anti-nausea medication was administered through the patient IV during the procedure. If the patient experiences nausea or vomiting after discharge, restrict diet to clear liquids (see above), until symptoms subside. If patient is experiencing persistent nausea or vomiting, please contact the ANESTHESIOLOGIST. EXPECTED ACTIVITY Patient may be tired and sleepy for the next several hours, and may take several naps. The patient should NOT drive, bike, swim, sign contracts, or engage in any other activity that requires full physical and mental coordination for the rest of the day. Patient may resume normal activities on the day following his/her surgery. FEVER Patient may experience a low-grade fever following anesthesia. Patient should stay indoors, and remain in a cool temperature controlled area. MEDICATIONS Please resume medications as prescribed unless otherwise indicated by the Anesthesiologist or Dentist. ADDITIONAL INSTRUCTIONS:	It is not uncommon for the patient to experience pain after his/her dental procedure. The anesthesiologist may have administered IV pain medications, and the dentist may have administered local anesthesia (numbing) to help the patient deal with potential pain. These medications typically wear off after a couple of hours. The following medications may be recommended for post-operative pain control. If the patient is experiencing			
Ibuprofen/Advil/Motrin (dose per package instructions) Immediately Begin atam/pm	, , , , , , , , , , , , , , , , , , ,	e per package instructions)		
NAUSEA & VOMITING Occasionally nausea and vomiting may occur following anesthesia. Anti-nausea medication was administered through the patient IV during the procedure. If the patient experiences nausea or vomiting after discharge, restrict diet to clear liquids (see above), until symptoms subside. If patient is experiencing persistent nausea or vomiting, please contact the ANESTHESIOLOGIST. EXPECTED ACTIVITY Patient may be tired and sleepy for the next several hours, and may take several naps. The patient should NOT drive, bike, swim, sign contracts, or engage in any other activity that requires full physical and mental coordination for the rest of the day. Patient may resume normal activities on the day following his/her surgery. FEVER Patient may experience a low-grade fever following anesthesia. Patient should stay indoors, and remain in a cool temperature controlled area. MEDICATIONS Please resume medications as prescribed unless otherwise indicated by the Anesthesiologist or Dentist. ADDITIONAL INSTRUCTIONS: I have reviewed these discharge instructions with my anesthesiologist and/or their assistant and have had all of my questions answered to	- OR -			
Occasionally nausea and vomiting may occur following anesthesia. Anti-nausea medication was administered through the patient IV during the procedure. If the patient experiences nausea or vomiting after discharge, restrict diet to clear liquids (see above), until symptoms subside. If patient is experiencing persistent nausea or vomiting, please contact the ANESTHESIOLOGIST. EXPECTED ACTIVITY Patient may be tired and sleepy for the next several hours, and may take several naps. The patient should NOT drive, bike, swim, sign contracts, or engage in any other activity that requires full physical and mental coordination for the rest of the day. Patient may resume normal activities on the day following his/her surgery. FEVER Patient may experience a low-grade fever following anesthesia. Patient should stay indoors, and remain in a cool temperature controlled area. MEDICATIONS Please resume medications as prescribed unless otherwise indicated by the Anesthesiologist or Dentist. ADDITIONAL INSTRUCTIONS: I have reviewed these discharge instructions with my anesthesiologist and/or their assistant and have had all of my questions answered to				
Patient may be tired and sleepy for the next several hours, and may take several naps. The patient should NOT drive, bike, swim, sign contracts, or engage in any other activity that requires full physical and mental coordination for the rest of the day. Patient may resume normal activities on the day following his/her surgery. FEVER Patient may experience a low-grade fever following anesthesia. Patient should stay indoors, and remain in a cool temperature controlled area. MEDICATIONS Please resume medications as prescribed unless otherwise indicated by the Anesthesiologist or Dentist. ADDITIONAL INSTRUCTIONS: I have reviewed these discharge instructions with my anesthesiologist and/or their assistant and have had all of my questions answered to	Occasionally nausea and vomiting may occur following anesthesia. Anti-nausea medication was administered through the patient IV during the procedure. If the patient experiences nausea or vomiting after discharge, restrict diet to clear liquids (see above), until symptoms subside. If patient is experiencing			
Patient may experience a low-grade fever following anesthesia. Patient should stay indoors, and remain in a cool temperature controlled area. MEDICATIONS Please resume medications as prescribed unless otherwise indicated by the Anesthesiologist or Dentist. ADDITIONAL INSTRUCTIONS: I have reviewed these discharge instructions with my anesthesiologist and/or their assistant and have had all of my questions answered to	Patient may be tired and sleepy for the next several hours, and may take several naps. The patient should NOT drive, bike, swim, sign contracts, or engage in any other activity that requires full physical and mental coordination for the rest of the day. Patient may resume normal activities on the day following his/her			
Please resume medications as prescribed unless otherwise indicated by the Anesthesiologist or Dentist. ADDITIONAL INSTRUCTIONS: I have reviewed these discharge instructions with my anesthesiologist and/or their assistant and have had all of my questions answered to	Patient may experience a low-grade fever following anesthesia. Patient should stay indoors, and remain in a			
I have reviewed these discharge instructions with my anesthesiologist and/or their assistant and have had all of my questions answered to				
I have reviewed these discharge instructions with my anesthesiologist and/or their assistant and have had all of my questions answered to satisfaction. I will receive a copy of these instructions and provide a contact number where I may be reached for the next 24 hours.	ADDITIONAL INSTRUCTIONS:			
Discharged to: Relationship:	Discharged to:	Relationship:		
Signature:Phone:	Signature:	Phone:		
Anesthesiologist: Phone: (800) 997-1220	Anesthesiologist:	Phone:(800) 997-1220		

Direct Line